

# Proteins with large influence on network dynamics evolve slowly

Ryan N. Gutenkunst

Center for Nonlinear Studies, Los Alamos National Laboratory, [ryang@lanl.gov](mailto:ryang@lanl.gov)


## Summary

Studies of protein evolution typically take a coarse view of protein function. Here I take a much finer view, using detailed mechanistic models to measure the influence of each constituent protein on its network's dynamics.

I show that a protein's dynamical influence and its evolutionary rate are negatively correlated, implying purifying selection on network dynamics. Correlation coefficients are typically of order -0.3, among the strongest known correlates of evolutionary rate. Moreover, this correlation is independent of interaction degree, expression level, and knock-out essentiality.

## Dynamical influence

Figure 1 - Dynamical influence



The inclusive measure of dynamics,  $\chi^2$ , sums over all dynamical species in the network [Gutenkunst (2007)].

$$\chi^2(\mathbf{k}) \propto \sum_y \int \left( \frac{y(t, \mathbf{k}) - y(t, \mathbf{k}^*)}{\sigma_y} \right)^2 dt$$

The dynamical influence  $\kappa_i$  of parameter  $i$  is (see Figure 1A)

$$\kappa_i = \sqrt{\frac{\partial^2 \chi^2}{\partial^2 \log k_i}}$$

The dynamical influence  $\mathcal{I}$  of a protein is the mean influence of parameters governing reactions the protein participates in.


$$\mathcal{I} = \langle \kappa_i \rangle_{\text{geom}}$$

For example, in the network shown in Figure 1B the influence of Ras incorporates all the highlighted reactions.

## Evolutionary rate


Maximum-likelihood dN/dS values were inferred with PAML, using alignments and sequences from Homologene and GenBank.

Figure 2 - Species Used



## Data

Figure 3 - Correlation in real networks



Knock-out essentiality in mouse:

●-essential ●-nonessential ●-no data

I tested all models in the BioModels database possessing 8 or more species annotated with Uniprot identifiers. Results are shown in Figure 3. As seen, for 9 out of 10 models there is a negative correlation between evolutionary rate and dynamical influence. Individual p-values are not dramatic, because model development is inherently low-throughput. However, it is compelling that so many systems show the expected correlation.

Note, the first 5 models all include an experimentally manipulated protein ligand, which I have removed from these analyses. I also omit the influence of multi-protein complexes, as it is not annotated which protein in a complex is active for each reaction. p-values are from one-sided permutation tests.

## Other potential correlates

It is known that protein evolutionary rates correlate with number of interaction partners (interaction degree), level of expression, and knock-out gene essentiality. I use the number of reactions  $R$  a protein is involved in (including complexes) as a proxy for interaction degree. I use the total amount  $X$  of a protein in the model as a proxy for expression. Essentiality in mouse  $K$  is measured using knock-out data from the Mouse Genome Database. Table 1 compares the correlations with and without controlling for these variables. It can be seen that these variables typically have little influence on the strength of correlation.

Table 1 - Correlations and partial correlations

Model	$r_{\mathcal{I},w}$ (p-val)	$r_{\mathcal{I},w,R}$	$r_{\mathcal{I},w,X}$	$r_{\mathcal{I},w,K}$
Birtwistle (2007)	-0.24 (0.20)	0.21	0.14	-0.31
Brown (2004)	-0.51 (0.04)	-0.38	-0.64	-0.56
Sasagawa (2005)	-0.45 (0.01)	-0.45	-0.46	0.21
Ung (2008)	-0.10 (0.32)	-0.15	-0.07	-0.47
Singh (2006)	-0.31 (0.16)	-0.35	-0.48	-0.36
Kim (2007)	-0.32 (0.17)	-0.34	-0.36	-0.25
Haberichter (2007)	-0.31 (0.19)	-0.50	-0.69	-0.10
Maeda (2006)	-0.20 (0.26)	-0.18	-0.22	-0.22
Yang (2007)	-0.53 (0.06)	-0.54	-0.65	-0.15
Neves (2008)	0.42 (0.87)	0.60	0.55	0.35

## References

- Birtwistle MR, Hatakeyama M, Yumoto N, Ogunnaik BA, Hoek JB, Kholodenko BN "Ligand-dependent responses of the ErbB signaling network: experimental and modeling analyses" *Mol Syst Biol* 3:144 (2007)
- Brown KS, Hill CC, Calero GA, Myers CR, Lee KH, Sethna JP, Cerione RA "The statistical mechanics of complex signaling networks: nerve growth factor signaling" *Phys Biol* 1:184 (2004)
- Gutenkunst RN, Waterfall JJ, Casey FP, Brown KS, Myers CR, Sethna JP "Universally sloppy parameter sensitivities in systems biology" *PLoS Comput Biol* 3(10):e189 (2007)
- Haberichter T, Madge B, Christopher RA, Yoshioka N, Dhiman A, Miller R, Gendelman R, Aksenov SV, Khalil IG, Dowdy SF "A systems biology dynamical model of mammalian G1 cell cycle progression" *Mol Syst Biol* 3:84 (2007)
- Kim D, Rath O, Kolch W, Cho KH "A hidden oncogenic positive feedback loop caused by crosstalk between Wnt and ERK Pathways" *Oncogene* 26:4571 (2007)
- Maeda A, Ozaki Y, Sivakumaran S, Akiyama T, Urakubo H, Usami A, Sato M, Kaibuchi K, Kuroda S "Ca-independent phospholipase A2-dependent sustained Rho-kinase activation exhibits all-or-none response" *Genes Cells* 11:1071 (2006)
- Neves SR, Tsokas P, Sarkar A, Grace EA, Rangamani P, Taubenfeld SM, Alberini CM, Schaff JC, Blitzer RD, Moraru II, Iyengar R "Cell shape and negative links in regulatory motifs together control spatial information flow in signaling networks" *Cell* 133:666 (2008)
- Sasagawa S, Ozaki Y, Fujita K, Kuroda S "Prediction and validation of the distinct dynamics of transient and sustained ERK activation" *Nat Cell Biol* 7:365 (2005)
- Singh A, Jayaraman A, Hahn J "Modeling regulatory mechanisms in IL-6 signal transduction in hepatocytes" *Biotechnol Bioeng* 95:850 (2006)
- Ung CY, Li H, Ma XH, Jia J, Li BW, Low BC, Chen YZ "Simulation of the regulation of EGFR endocytosis and EGFR-ERK signaling by endophilin-mediated RhoA-EGFR crosstalk" *FEBS Lett* 582:2283 (2008)
- Yang K, Ma W, Liang H, Ouyang Q, Tang C, Lai L "Dynamic simulations on the arachidonic acid metabolic network" *PLoS Comput Biol* 3:e55 (2007)